arXiv:math-ph/9902018v1 15 Feb 1999

A note on Farey sequences and Hausdorff dimension

Wellington da Cruz*
Departamento de Física,
Universidade Estadual de Londrina, Caixa Postal 6001,
Cep 86051-990 Londrina, PR, Brazil

(February 7, 2008)

Abstract

We prove that the Farey sequences can be express into equivalence classes labeled by a fractal parameter which looks like a Hausdorff dimension h defined within the interval $1 < h < 2$. The classes h satisfy the same properties of the Farey series and for each value of h there exists an algebraic equation.

PACS numbers: 02.10.Lh, 02.30.Lt
Keywords: Farey sequences; Hausdorff dimension; Number theory

Typeset using REVTEX

---

*E-mail: wdacruz@exatas.campus.uel.br

From considerations about Fractional Quantum Hall Effect (FQHE) [1] we have found a connection between a fractal parameter or Hausdorff dimension h and the Farey series [2]. Thus, we have the following **theorem**:

The elements of the Farey series belong to distinct equivalence classes labeled by a fractal parameter h defined into the interval $1 < h < 2$, such that these classes satisfy the same properties observed for that fractions. Also, for each value of h there exists an algebraic equation .

The fractal parameter h is related to ν (an irreducible number $\frac{p}{q}$, with p and q integers) as follows

$$\begin{array}{llll} h-1=1-\nu, & 0<\nu<1; & h-1=\nu-1, & 1<\nu<2; \\ h-1=3-\nu, & 2<\nu<3; & h-1=\nu-3, & 3<\nu<4; \\ h-1=5-\nu, & 4<\nu<5; & h-1=\nu-5, & 5<\nu<6; \\ h-1=7-\nu, & 6<\nu<7; & h-1=\nu-7, & 7<\nu<8; \\ h-1=9-\nu, & 8<\nu<9; & h-1=\nu-9, & 9<\nu<10; \\ etc, & & & \end{array} \tag{1}$$

We can extract, for example, the classes

$$\begin{array}{ll} \left\{\frac{1}{3},\frac{5}{3},\frac{7}{3},\frac{11}{3},\cdots\right\}_{h=\frac{5}{3}}, & \left\{\frac{5}{14},\frac{23}{14},\frac{33}{14},\frac{51}{14},\cdots\right\}_{h=\frac{23}{14}}; \\ \left\{\frac{4}{11},\frac{18}{11},\frac{26}{11},\frac{40}{11},\cdots\right\}_{h=\frac{18}{11}}, & \left\{\frac{7}{19},\frac{31}{19},\frac{45}{19},\frac{69}{19},\cdots\right\}_{h=\frac{31}{19}}; \\ \left\{\frac{10}{27},\frac{44}{27},\frac{64}{27},\frac{98}{27},\cdots\right\}_{h=\frac{44}{27}}, & \left\{\frac{3}{8},\frac{13}{8},\frac{19}{8},\frac{29}{8},\cdots\right\}_{h=\frac{13}{8}}; \\ \left\{\frac{3}{7},\frac{11}{7},\frac{17}{7},\frac{25}{7},\cdots\right\}_{h=\frac{11}{7}}, & \left\{\frac{4}{9},\frac{14}{9},\frac{22}{9},\frac{32}{9},\cdots\right\}_{h=\frac{14}{9}} \\ \left\{\frac{5}{11},\frac{17}{11},\frac{27}{11},\frac{39}{11},\cdots\right\}_{h=\frac{17}{11}}, & \left\{\frac{6}{13},\frac{20}{13},\frac{32}{13},\frac{46}{13},\cdots\right\}_{h=\frac{20}{13}}, \\ \left\{\frac{2}{5},\frac{8}{5},\frac{12}{5},\frac{18}{5},\cdots\right\}_{h=\frac{8}{5}}, & \end{array} \tag{2}$$

and for that we can consider the serie (h,ν)

$$\begin{array}{l} \left(\frac{5}{3},\frac{1}{3}\right)\rightarrow\left(\frac{18}{11},\frac{4}{11}\right)\rightarrow\left(\frac{13}{8},\frac{3}{8}\right)\rightarrow\left(\frac{8}{5},\frac{2}{5}\right)\rightarrow \\ \left(\frac{11}{7},\frac{3}{7}\right)\rightarrow\left(\frac{14}{9},\frac{4}{9}\right)\rightarrow\left(\frac{17}{11},\frac{5}{11}\right)\rightarrow\left(\frac{20}{13},\frac{6}{13}\right)\rightarrow\cdots \end{array} \tag{3}$$

The classes h satisfy all properties of the Farey series:

P1. If $h_1=\frac{p_1}{q_1}$ and $h_2=\frac{p_2}{q_2}$ are two consecutive fractions $\frac{p_1}{q_1}>\frac{p_2}{q_2}$, then $|p_2q_1-q_2p_1|=1$.

P2. If $\frac{p_1}{q_1}$, $\frac{p_2}{q_2}$, $\frac{p_3}{q_3}$ are three consecutive fractions $\frac{p_1}{q_1}>\frac{p_2}{q_2}>\frac{p_3}{q_3}$, then $\frac{p_2}{q_2}=\frac{p_1+p_3}{q_1+q_3}$.

P3. If $\frac{p_1}{q_1}$ and $\frac{p_2}{q_2}$ are consecutive fractions in the same sequence, then among all fractions between the two, $\frac{p_1+p_2}{q_1+q_2}$ is the unique reduced fraction with the smallest denominator.

For more details about Farey series see [2]. All these properties can be verified for the classes considered above as an example. Another example is

$$\begin{aligned}(h,\nu)=\left(\frac{11}{6},\frac{1}{6}\right)&\rightarrow\left(\frac{9}{5},\frac{1}{5}\right)\rightarrow\left(\frac{7}{4},\frac{1}{4}\right)\rightarrow\left(\frac{5}{3},\frac{1}{3}\right)\rightarrow\\ &\left(\frac{8}{5},\frac{2}{5}\right)\rightarrow\left(\frac{3}{2},\frac{1}{2}\right)\rightarrow\left(\frac{7}{5},\frac{3}{5}\right)\rightarrow\left(\frac{4}{3},\frac{2}{3}\right)\rightarrow\\ &\left(\frac{5}{4},\frac{3}{4}\right)\rightarrow\left(\frac{6}{5},\frac{4}{5}\right)\rightarrow\left(\frac{7}{6},\frac{5}{6}\right)\rightarrow\cdots,\end{aligned} \tag{4}$$

where the ν sequence is the Farey series of order 6. Thus, we observe that because of the *fractal spectrum* (Eq.1), we can write down any Farey series of rational numbers.

In summary, we have extracted from considerations about FQHE [1] a beautiful connection between number theory and physics. We have shown that the Farey series can be arranged into equivalence classes labeled by a fractal parameter h which looks like a Hausdorff dimension. For each value of h we have an algebraic equation derived of the functional equation

$$\xi=\{\mathcal{Y}(\xi)-1\}^{h-1}\{\mathcal{Y}(\xi)-2\}^{2-h}, \tag{5}$$

where ξ is an exponential function. Then, there exists a relation between algebraic equation and Farey series. The connection between a geometric parameter related to paths of particles (charge-flux system called *fractons* [1,3] identified with holes of a multiply connected space) defined in the context of a two dimensional physical system and rational numbers deserve a more deep research.

REFERENCES

[1] W. da Cruz, preprint/UEL-DF/981003, cond-mat/9810381.
[2] M. Schroeder, *Number Theory in Science and Communication* (Springer Verlag, 1997)
[3] W. da Cruz, preprint/UEL-DF/981004, cond-mat/9811007.